


BRITISH GEOLOGICAL SURVEY

Lerwick Observatory

Monthly Magnetic Bulletin

September 2011

11/09/LE



SHETLAND ISLANDS



British Geological Survey

NATURAL ENVIRONMENT RESEARCH COUNCIL

LERWICK OBSERVATORY MAGNETIC DATA

1. Introduction

Lerwick observatory is one of three geomagnetic observatories in the UK operated and maintained by the British Geological Survey (BGS).

This bulletin is published to provide rapid access to the provisional geomagnetic observatory results. The information is freely available for personal, academic, educational and non-commercial research or use. Magnetic observatory data are presented as a series of plots of one-minute, hourly and daily values, followed by tabulations of monthly values, reports of rapid variations and geomagnetic activity indices. The operation of the observatory and presentation of data are described in the rest of this section.

Enquiries about the data should be addressed to:

Geomagnetism Team
Earth Hazards and Systems
British Geological Survey
Murchison House, West Mains Road
Edinburgh EH9 3LA
Scotland, UK

Tel: +44 (0) 131 667 1000
Fax: +44 (0) 131 650 0265
E-mail: enquiries@bgs.ac.uk
Internet: www.geomag.bgs.ac.uk

2. Position

The observatory is situated on a ridge of high ground about 2.5 km to the SW of the port of Lerwick in Shetland. The observatory co-ordinates are:

Geographic: 60.133°N 358.816°E
Geomagnetic: 61.876°N 88.858°E
Height above mean sea level: 85 m

The geomagnetic co-ordinates are approximations, calculated using the 11th generation International Geomagnetic Reference Field (IGRF) at epoch 2011.5. On-line access to models (including IGRF), charts and navigational data are available at www.geomag.bgs.ac.uk/data_service/models_compass/home

3. The Observatory Operation

3.1 GDAS

The observatory operates under the control of the Geomagnetic Data Acquisition System (GDAS),

which was developed by BGS staff, installed in 2002, and became fully operational in January 2003. The data acquisition software, running on QNX operated computers, controls the data logging and the communications.

There are two sets of sensors used for making magnetic measurements. A tri-axial linear-core fluxgate magnetometer, manufactured by the Danish Meteorological Institute, is used to measure the variations in the horizontal (H) and vertical (Z) components of the field. The third sensor is oriented perpendicular to these, and measures variations, which are proportional to the changes in declination (D). Measurements are made at a rate of 1 Hz.

In addition to the fluxgate sensors there is a proton precession magnetometer (PPM) making measurements of the absolute total field intensity (F) at a rate of 0.1Hz.

The raw unfiltered data are retrieved automatically via Internet connections to the BGS office in Edinburgh in near real-time. The fluxgate data are filtered to produce one-minute values using a 61-point cosine filter and the total field intensity samples are filtered using a 7-point cosine filter. The one-minute values provide input for various data products, available on-line at www.geomag.bgs.ac.uk/data_service/home

3.2 Back-up Systems

There are two other fully independent identical systems, GDAS 2 and GDAS 3, operating at the observatory. The data from these are also processed in near real-time and used for quality control purposes. They are also used to fill any gaps or replace any corrupt values in the primary system, GDAS 1.

3.3 Absolute Observations

The GDAS fluxgate magnetometers accurately measure variations in the components of the geomagnetic field, but not the absolute magnitudes. Two sets of absolute measurements of the field are made manually once per week. A fluxgate sensor mounted on a theodolite is used to determine D and inclination (I); the GDAS PPM measurements, with a site difference correction applied, are used for F . The absolute observations are used in conjunction with the GDAS variometer measurements to produce a continuous record of

the absolute values of the geomagnetic field elements as if they had been measured at the observatory reference pillar.

4. Observatory Results

The data presented in the bulletin are in the form of plots and tabulations described in the following sections.

4.1 Absolute Observations

The absolute observation measurements made during the month are tabulated. Also included are the corresponding baseline values, which are the differences between the absolute measurements and the variometer measurements of D , H and Z (in the sense absolute–variometer). These are also plotted (markers) along with the derived preliminary daily baseline values (line) throughout the year. Daily mean differences between the measured absolute F and the F computed from the baseline corrected H and Z values are plotted in the fourth panel (in the sense measured–derived). The bottom panel shows the daily mean temperature in the fluxgate chamber.

4.2 Summary magnetograms

Small-scale magnetograms are plotted which allow the month's data to be viewed at a glance. They are plotted 16 days to a page and show the one-minute variations in D , H and Z . The scales are shown on the right-hand side of the page. On disturbed days the scales are multiplied by a factor, which is indicated above the panel for that day. The variations are centred on the monthly mean value, shown on the left side of the page.

4.3 Magnetograms

The daily magnetograms are plotted using one-minute values of D , H and Z from the fluxgate sensors, with any gaps filled using back-up data. The magnetograms are plotted to a variable scale; scale bars are shown to the right of each plot. The absolute level (the monthly mean value) is indicated on the left side of the plots.

4.4 Hourly Mean Value Plots

Hourly mean values of D , H and Z for the past 12 months are plotted in 27-day segments corresponding to the Bartels solar rotation number. Magnetic disturbances associated with active regions and/or coronal holes on the Sun may recur after 27 days: the same is true for geomagnetically quiet intervals. Plotting the data in this way highlights this recurrence. Diurnal variations are also clear in these plots and the amplitude changes

throughout the year highlight the seasonal changes. Longer term secular variation is also illustrated.

Full lists of the UK observatory hourly mean values from 1983 to the present day are available at www.geomag.bgs.ac.uk/data_service/data/obs_data/hourly_means

4.5 Daily and Monthly Mean Values

Daily mean values of D , H , Z and F are plotted throughout the year. In addition, a table of monthly mean values of all the geomagnetic elements is provided. These values depend on accurate specification of the fluxgate sensor baselines. It is anticipated that these provisional values will not be altered by more than a few nT or tenths of arcminutes before being made definitive at the end of the year.

4.6 Rapid Variations

Charged particles stream from the Sun in the solar wind. The solar wind interacts with the geomagnetic field to create a cavity, the magnetosphere, in which the field is confined. When a region of enhanced velocity and/or density in the solar wind arrives at the dayside boundary of the magnetosphere (at about 10 earth radii) the boundary is pushed towards the Earth. Currents set up on the boundary of the magnetosphere can cause an abrupt change in the geomagnetic field measured on the ground and this is recorded on observatory magnetograms as a sudden impulse (*si*). If, following an *si*, there is a change in the rhythm of activity, the *si* is termed a storm sudden commencement (*ssc*). A classical magnetic storm exhibiting initial, main and recovery phases (shown by, for instance, the *Dst* ring current index) can often occur after a *ssc*, in which case the start of the storm is taken as the time of the *ssc*.

Solar flares, seen at optical wavelengths as a sudden brightening of a small region of the Sun's surface, are also responsible for increased X-ray emissions. These X-rays cause increased ionisation in the ionosphere, which leads to absorption of short-wave radio signals. A solar flare effect (*sfe*), or "crochet", may be observed on a magnetogram during geomagnetically quiet times. It is a relatively short-term change (tens of minutes) to the normal diurnal variation and can vary in size (tens of nT) depending on local time (LT), geomagnetic latitude and solar zenith angle.

4.7 Local geomagnetic activity indices

The Observatory K index. This summarises geomagnetic activity at an observatory by assigning a code, an integer in the range 0 to 9, to each 3-hour Universal Time (UT) interval. The index for each 3-hour UT interval is determined from the maximum range in H or D (scaled in nT), with allowance made for the regular (undisturbed) diurnal variation. The conversion from range to an index value is made using a quasi-logarithmic scale, with the scale values dependent on the geomagnetic latitude of the observatory. The lower bounds (in nT) for the classification of each period at Lerwick are:

0	1	2	3	4	5	6	7	8	9
0	10	20	40	80	140	240	400	660	1000

The K index retains the LT and seasonal dependence of activity associated with the position of the observatory. The 3-hourly K indices for the month are tabulated and also plotted as a histogram. All UK observatory K indices are available at

www.geomag.bgs.ac.uk/data_service/data/magnetic_indices/k_indices

4.8 Global geomagnetic activity indices

The aa index. A number of 3-hour geomagnetic indices are computed by combining K indices from networks of observatories to characterise global activity levels and to eliminate LT and seasonal effects. The simplest of these is the aa index, computed using the K indices from two approximately antipodal observatories: Hartland in the UK and Canberra in Australia. The aa index is calculated from linearisations of the Hartland and Canberra K indices, and has units of nT. The 3-hourly aa indices are tabulated along with the daily mean value of aa (denoted Aa), the mean values of aa for the intervals 00-12UT (Aa_{am}) and 12-24UT (Aa_{pm}) and the monthly mean value. The 3-hourly aa indices for the month are also plotted as a histogram.

Although the aa index is based on data from only two observatories, provided averages over 12 hours or longer are used, the index is strongly correlated with the ap and am indices, which are derived using data from more extensive observatory networks.

The aa indices listed in this bulletin are available at www.geomag.bgs.ac.uk/data_service/data/magnetic_indices/aaindex as well as the full data set from 1868.

Definitive aa are published by the International Service for Geomagnetic Indices, LATMOS, 4 Avenue de Neptune, F-94107 Saint Maur Cedex, France.

5. Conditions of Use

The data presented in this bulletin are provided for personal, academic, educational, non-commercial research or other non-commercial use and are not for sale or distribution to third parties without written permission from BGS.

Reproduction of any part of this bulletin should be accompanied by the statement: 'Reproduced with the permission of the British Geological Survey ©NERC. All rights Reserved'. Publications making use of the data should include an acknowledgment statement of the form: 'The results presented in this paper rely on the data collected at Lerwick magnetic observatory, operated by the British Geological Survey.'

Commercial users can contact the geomagnetism team for information on the range of applications and services offered. Full contact details are available at www.geomag.bgs.ac.uk/contactus/staff

**This product includes mapping data licensed from Ordnance Survey with the permission of HMSO © Crown copyright. All rights reserved.
Licence Number: 100017897/2011**

LERWICK OBSERVATORY

ABSOLUTE OBSERVATIONS

Date	Day Number	Declination			Inclination		Total Field		Horizontal Intensity		Vertical Intensity		Observer
		Time (UT)	Absolute (°)	Baseline (°)	Time (UT)	Absolute (°)	Site difference (nT)	Absolute corrected (nT)	Absolute (nT)	Baseline (nT)	Absolute (nT)	Baseline (nT)	
05-Sep-11	248	08:29	-2.5837	-3.0233	08:40	72.8164	-0.2	50795.7	15006.8	14965.2	48528.3	48417.1	PK
05-Sep-11	248	08:50	-2.5992	-3.0383	09:03	72.8154	-0.2	50796.1	15007.8	14966.6	48528.4	48416.7	PK
12-Sep-11	255	08:36	-2.6369	-3.0183	08:51	72.8361	-0.2	50768.0	14981.9	14966.9	48507.0	48416.7	WH
12-Sep-11	255	09:03	-2.6651	-3.0350	09:17	72.8407	-0.2	50767.4	14977.9	14966.0	48507.6	48417.0	WH
19-Sep-11	262	08:31	-2.5590	-3.0117	08:47	72.8442	-0.2	50799.7	14984.4	14967.0	48539.4	48416.7	PD
19-Sep-11	262	08:58	-2.5645	-3.0117	09:09	72.8518	-0.2	50796.3	14977.0	14963.6	48538.1	48417.8	PD
21-Sep-11	264	16:12	-2.7033	-3.0167	16:59	72.8163	-0.2	50809.6	15011.0	14962.6	48541.6	48418.2	NL
23-Sep-11	266	08:28	-2.5888	-3.0250	08:45	72.8130	-0.2	50791.6	15008.5	14969.7	48523.5	48416.0	WH
23-Sep-11	266	08:58	-2.5995	-3.0267	09:14	72.8178	-0.2	50790.5	15004.1	14968.5	48523.8	48416.5	WH
26-Sep-11	269	08:12	-2.5695	-3.0233	08:25	72.8101	-0.2	50793.6	15011.5	14967.6	48524.7	48416.6	PD
26-Sep-11	269	08:35	-2.5731	-3.0217	08:46	72.8147	-0.2	50791.6	15007.0	14967.3	48524.0	48416.7	PD

Lerwick 2011

Declination : Absolute - Variometer (markers) and Baseline Applied (line)

-181.5

5min

Horizontal Intensity : Absolute - Variometer (markers) and Baseline Applied (line)

14968

20nT

Vertical Intensity : Absolute - Variometer (markers) and Baseline Applied (line)

48417

20nT

Total Intensity : Absolute (Proton F) - Baseline Adjusted Variometer (derived F)

0.0


20nT

Temperature Inside Variometer Hut

18.2

10°C


Jan Feb Mar Apr May Jun Jul Aug Sep Oct Nov Dec




Lerwick

September

2011





Lerwick


September

2011

Date: 01-09-2011


Lerwick

Day number: 244



Date: 02-09-2011


Day number: 245



Date: 03-09-2011


Lerwick

Day number: 246



Date: 04-09-2011


Day number: 247



Date: 05-09-2011


Lerwick

Day number: 248



Date: 06-09-2011


Day number: 249



Date: 07-09-2011


Lerwick

Day number: 250



Date: 08-09-2011


Day number: 251



Date: 09-09-2011


Lerwick

Day number: 252



Date: 10-09-2011


Day number: 253



Date: 11-09-2011


Lerwick

Day number: 254



Date: 12-09-2011


Day number: 255



Date: 13-09-2011


Lerwick

Day number: 256



Date: 14-09-2011


Day number: 257



Date: 15-09-2011


Lerwick

Day number: 258



Date: 16-09-2011


Day number: 259



Date: 17-09-2011


Lerwick

Day number: 260



Date: 18-09-2011


Day number: 261



Date: 19-09-2011


Lerwick

Day number: 262



Date: 20-09-2011


Day number: 263



Date: 21-09-2011


Lerwick

Day number: 264



Date: 22-09-2011


Day number: 265



Date: 23-09-2011


Lerwick

Day number: 266



Date: 24-09-2011


Day number: 267



Date: 25-09-2011


Lerwick

Day number: 268



Date: 26-09-2011


Day number: 269



Date: 27-09-2011


Lerwick

Day number: 270



Date: 28-09-2011


Day number: 271



Date: 29-09-2011


Lerwick

Day number: 272




Date: 30-09-2011

Day number: 273




Lerwick Observatory: Declination (degrees)




Hourly Mean Values Plotted By Bartels Solar Rotation Number


Lerwick Observatory: Horizontal Intensity (nT)



Lerwick Observatory: Vertical Intensity (nT)



Lerwick Observatory 2011



Monthly Mean Values for Lerwick Observatory 2011

Month	<i>D</i>	<i>H</i>	<i>I</i>	<i>X</i>	<i>Y</i>	<i>Z</i>	<i>F</i>
January	-2° 47.4′	15008 nT	72° 48.6′	14991 nT	-730 nT	48513 nT	50782 nT
February	-2° 46.2′	15006 nT	72° 48.8′	14988 nT	-725 nT	48517 nT	50784 nT
March	-2° 45.1′	15007 nT	72° 48.8′	14989 nT	-720 nT	48516 nT	50784 nT
April	-2° 44.1′	15008 nT	72° 48.6′	14991 nT	-716 nT	48516 nT	50785 nT
May	-2° 43.2′	15013 nT	72° 48.3′	14996 nT	-712 nT	48516 nT	50786 nT
June	-2° 42.0′	15017 nT	72° 48.2′	15000 nT	-708 nT	48521 nT	50791 nT
July	-2° 41.0′	15017 nT	72° 48.2′	15000 nT	-703 nT	48520 nT	50791 nT
August	-2° 39.7′	15010 nT	72° 48.7′	14994 nT	-697 nT	48524 nT	50792 nT
September	-2° 38.5′	15004 nT	72° 49.1′	14988 nT	-692 nT	48527 nT	50793 nT

Note

i. The values shown here are provisional.

LERWICK RAPID VARIATIONS

SI and SSCs

Date	Time (UT)	Type	Quality	H (nT)	D (min)	Z (nT)
09-09-11	12 42	SSC*	A	57.7	-7.73	21.8
17-09-11	03 43	SSC*	B	6.1	-4.11	1.4/1.8
25-09-11	11 45	SI*	B	-12.1	1.62	-3.7
26-09-11	12 35	SSC*	A	41.9	-8.13	15.3

Notes:

An asterisk (*) indicates that the principal impulse was preceded by a smaller reversed impulse.

The quality of the event is classified as follows:

A = very distinct

B = fair, ordinary, but unmistakable

C = doubtful

The amplitudes given are for the first chief movement of the event.

SFEs

Date	Universal Time			H (nT)	D (min)	Z (nT)
	Start	Maximum	End			
24-09-11	09 35	09 41	09 59	-9.7	1.92	2.0


Note:

The amplitudes given are for the first chief movement of the event.

INDICES OF GEOMAGNETIC ACTIVITY

Day	K - INDICES FOR THREE-HOUR INTERVAL							
	00-03	03-06	06-09	09-12	12-15	15-18	18-21	21-24
1	1	0	0	0	1	1	0	1
2	0	1	0	0	1	1	1	1
3	2	2	2	2	4	3	2	2
4	2	3	1	0	2	3	2	0
5	1	0	0	0	1	3	3	1
6	1	2	2	1	2	1	1	1
7	3	1	0	1	1	1	2	0
8	0	1	0	0	1	0	0	0
9	0	0	1	1	4	6	5	6
10	6	4	2	4	2	4	4	5
11	3	1	0	1	3	2	2	4
12	4	3	3	2	3	3	4	4
13	4	4	3	2	2	1	2	2
14	0	0	1	1	1	1	1	1
15	2	0	1	1	2	2	2	1
16	0	0	0	0	1	1	0	1
17	1	2	3	4	4	4	4	3
18	1	0	0	1	1	1	0	1
19	0	0	0	0	0	1	1	0
20	0	2	0	1	1	1	1	2
21	0	0	0	1	2	2	0	1
22	1	0	0	1	1	1	1	2
23	1	0	0	0	0	0	0	0
24	0	0	0	1	1	0	1	1
25	0	0	0	2	1	2	1	0
26	1	0	1	1	3	7	6	6
27	5	4	3	2	3	2	2	4
28	4	3	3	1	3	3	2	1
29	4	3	2	1	2	3	2	1
30	2	0	0	0	0	0	2	1

Lerwick Observatory 3-hourly K-Indices



The *aa* Index

Date	Day	3-hourly <i>aa</i> -indices								<i>Aa_{am}</i>	<i>Aa_{pm}</i>	<i>Aa</i>
01-09-11	244	5	8	2	2	5	5	2	5	4.5	4.5	4.5
02-09-11	245	2	9	5	9	12	5	5	8	6.5	7.5	7.0
03-09-11	246	16	32	16	32	24	32	9	12	24.1	19.3	21.7
04-09-11	247	24	32	12	5	24	45	20	8	18.3	24.3	21.3
05-09-11	248	8	2	5	8	12	24	32	8	5.8	19.0	12.4
06-09-11	249	5	12	24	12	16	16	12	8	13.3	13.0	13.1
07-09-11	250	20	5	5	12	12	5	17	5	10.5	9.9	10.2
08-09-11	251	5	8	2	2	2	5	2	2	4.5	3.1	3.8
09-09-11	252	2	8	5	8	81	80	102	115	5.8	94.3	50.0
10-09-11	253	80	45	46	80	24	67	67	67	62.8	55.9	59.4
11-09-11	254	24	16	8	8	46	16	12	45	13.9	29.8	21.8
12-09-11	255	45	24	46	46	32	46	80	67	40.2	56.1	48.2
13-09-11	256	45	59	45	38	16	5	12	12	46.9	11.2	29.1
14-09-11	257	8	5	8	12	8	8	8	12	8.2	8.8	8.5
15-09-11	258	9	12	8	8	12	16	12	12	9.2	12.9	11.1
16-09-11	259	2	2	5	8	5	8	5	5	4.5	5.8	5.1
17-09-11	260	5	24	59	102	80	102	45	37	47.5	66.1	56.8
18-09-11	261	12	12	2	16	16	8	5	16	10.6	11.3	10.9
19-09-11	262	2	2	8	5	5	5	8	2	4.5	5.1	4.8
20-09-11	263	9	24	16	16	16	5	8	9	16.4	9.5	13.0
21-09-11	264	2	2	8	12	32	12	5	16	6.2	16.3	11.2
22-09-11	265	5	2	5	5	8	12	16	12	4.5	11.9	8.2
23-09-11	266	5	5	2	2	2	5	2	12	3.8	5.5	4.7
24-09-11	267	9	2	5	16	12	5	12	12	8.2	10.2	9.2
25-09-11	268	5	9	5	16	12	16	5	8	8.9	10.2	9.6
26-09-11	269	12	5	8	12	81	136	171	80	9.2	117.1	63.2
27-09-11	270	59	59	46	32	32	24	32	59	48.9	36.8	42.9
28-09-11	271	45	59	46	16	46	24	12	12	41.5	23.4	32.5
29-09-11	272	137	116	46	16	32	24	24	12	78.7	23.0	50.9
30-09-11	273	24	5	2	2	2	8	24	32	8.5	16.6	12.6
Monthly Mean Value											21.9	

Notes

- i. The units of the *aa* index are nT.
- ii. The 3-hour *aa* values are rounded to the nearest integer. Where $aa = *.5$, *aa* is rounded down.
- iii. Daily values (Aa_{am} , Aa_{pm} and *Aa*) are computed from *aa* values of original resolution.
- iv. The monthly mean value is computed from the daily mean values, *Aa*.
- v. Definitive *aa* indices are derived and published by the International Service for Geomagnetic Indices.

3-hourly *aa*-indices

